

FRÉCHET-URYSOHN FANS IN FREE TOPOLOGICAL GROUPS

Taras Banakh, Dušan Repovš, and Lyubomyr Zdomskyy

November 24, 2007

Abstract

In this paper we answer the question of T. Banakh and M. Zarichnyi constructing a copy of the Fréchet-Urysohn fan S_ω in a topological group G admitting a functorial embedding $[0, 1] \subset G$. The latter means that each autohomeomorphism of $[0, 1]$ extends to a continuous homomorphism of G . This implies that many natural free topological group constructions (e.g. the constructions of the Markov free topological group, free abelian topological group, free totally bounded group, free compact group) applied to a Tychonov space X containing a topological copy of the space \mathbb{Q} of rationals give topological groups containing S_ω .

1. Introduction

D. Shakhmatov noticed in [16] that the classical Lefschetz-Nöbeling-Pontryagin Theorem on embeddings of n -dimensional compacta into \mathbb{R}^{2n+1} has no categorical counterpart: one cannot embed every finite-dimensional compact space X into a finite-dimensional topological group FX so that each continuous map $f : X \rightarrow Y$ extends to a continuous group homomorphism $Ff : FX \rightarrow FY$. The proof of this fact exploited Kulesza's example of a pathological 1-dimensional (non-metrizable) compact space that cannot be embedded into a finite-dimensional topological group [11]. However, it was discovered in [4] that the problem lies already at the level of the unit interval $[0, 1]$, since it admits no functorial embedding into any metrizable or finite-dimensional group. So, each topological group containing a functorially embedded interval is non-metrizable and thus has uncountable character.

In light of this let us remark that the Markov free topological group $F_M I$ over the interval $I = [0, 1]$ has character $\chi(F_M I) = \mathfrak{d}$ (see [14], [15]), where \mathfrak{d} is the well-known uncountable small cardinal equal to the cofinality of the poset (ω^ω, \leq) . This cardinal is equal to the cardinality of continuum \mathfrak{c} under Martin's Axiom, but can also be strictly smaller than \mathfrak{c} in some models of ZFC (see [6], [18]).

In this paper we show that the inequality $\chi(FI) \geq \mathfrak{d}$ holds for many other free topological group constructions. First we give precise definitions.

Let \mathcal{T} be a subcategory of the category \mathcal{Top} of topological spaces and their continuous maps. By a *functor of a free topological group* on \mathcal{T} we understand a pair (F, i) consisting of

⁰The authors were supported by research grants P1-0292-0101-04 and BI-UA/04-06-007. We thank the referees and the Editor for several comments and suggestions.

2000 Math. Subj. Classif.: 18B30, 22A05, 54C20, 54E35, 54F15.

Keywords and phrases. Free topological group, functorial embedding, sequential fan, small cardinals.

a covariant functor $F : \mathcal{T} \rightarrow \mathcal{G}$ from \mathcal{T} into the category \mathcal{G} of topological groups and their continuous homomorphisms, and a natural transformation $i = \{i_X\} : \text{Id} \rightarrow F$ of the identity functor $\text{Id} : \mathcal{T} \rightarrow \mathcal{T}$ into the functor F whose components $i_X : X \rightarrow FX$ are topological embeddings for all spaces $X \in \mathcal{T}$. The naturality of i means that for any morphism $f : X \rightarrow Y$ in \mathcal{T} we have the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{i_X} & FX \\ f \downarrow & & \downarrow Ff \\ Y & \xrightarrow{i_Y} & FY \end{array}$$

Therefore a functor of a free topological group (F, i) to any topological space $X \in \text{Ob}(\mathcal{T})$ assigns a topological group FX containing a topological copy $i_X(X) \subset FX$ of X so that each morphism $f : X \rightarrow Y$ to another object of \mathcal{T} extends in a canonical way to a group homomorphism $Ff : FX \rightarrow FY$. A functor (F, i) is said to be *minimal*, if for every space $X \in \text{Ob}(\mathcal{T})$ the group FX is algebraically generated by $i_X(X)$. In the case when \mathcal{T} has only one object X and $\text{Mor}(X, X)$ coincides with the set of all autohomeomorphisms of X , the embedding $i_X : X \rightarrow FX$ is simply called [4] a *functorial embedding* of X into the group $G = FX$. It was proven in [4] that if there exists a functorial embedding of the interval $I = [0, 1]$ into a topological group G , then G is non-metrizable and infinite-dimensional. In this paper we shall make this result more precise by showing that such a group G contains a topological copy of the quotient space $IS_\omega = [0, 1] \times \omega / \{0\} \times \omega$, called the *sequential hedgehog* by analogy with the sequential (or alternatively Fréchet-Urysohn) fan $S_\omega = S_0 \times \omega / \{0\} \times \omega$, where $S_0 = \{0\} \cup \{1/n : n > 0\}$ is the convergent sequence. This answers a question stated in [4], and implies that $\chi(G) \geq \mathfrak{d}$.

Theorem 1. *If there exists a functorial embedding of the closed interval $[0, 1]$ into a topological group G , then G contains a copy of the sequential hedgehog IS_ω .*

A subcategory \mathcal{T} of the category \mathcal{Top} of topological spaces is said to be *full* if any continuous map between objects of \mathcal{T} is a morphism in \mathcal{T} .

Theorem 2. *Let (F, i) be a functor of a free topological group on a full subcategory \mathcal{T} of \mathcal{Top} , containing the interval $I = [0, 1]$ as an object. For every space $X \in \text{Ob}(\mathcal{T})$ containing a copy of \mathbb{Q} (resp. I), the topological group FX contains a copy of the Fréchet-Urysohn fan S_ω (resp. sequential hedgehog IS_ω) and hence has character $\chi(FX) \geq \mathfrak{d}$.*

It is interesting to remark that Theorem 2 is not true for the category $\mathcal{T} = \mathcal{C}(0)$ of zero-dimensional compacta and their continuous maps. For example, the functor F which assigns to each zero-dimensional compact space X the compact group $\mathbb{Z}_2^{\text{Mor}(X, \mathbb{Z}_2)}$ containing a diagonally embedded copy of X , is a functor of a free topological group (see [16]). For any compact metrizable space X the group $\mathbb{Z}_2^{\text{Mor}(X, \mathbb{Z}_2)}$ is metrizable and hence contains no copy of the Fréchet-Urysohn fan. This shows that the inclusion $I \in \text{Ob}(\mathcal{T})$ in Theorem 2 is essential.

Problem 3. *Let (F, i) be a functor of a free topological group on a full subcategory \mathcal{I} of \mathcal{Top} . Assume that $I \in \text{Ob}(\mathcal{T})$ and X is an object of \mathcal{T} . Does FX contain a copy of S_ω if X contains a non-trivial convergent sequence? (This is true for the functor of the Markov free topological group).*

We recall that a topological space X is said to be *scattered*, if every non-empty subspace Y of X has an isolated point. Combining Theorem 2 with the main result of [3], we shall derive the following

Corollary 1. *Let (F, i) be a minimal functor of a free topological group on a full category \mathcal{T} of topological spaces such that $I \in \text{Ob}(\mathcal{T})$. Suppose that $X \in \text{Ob}(\mathcal{T})$ is a metrizable separable space that has a compactification $\tilde{X} \in \text{Ob}(\mathcal{T})$. If FX is a k -space, then X is locally compact or scattered.*

This corollary can be compared with a result of Arkhangel'skiĭ, Okunev and Pestov [2] who proved that for a metrizable space X the Markov free topological group $F_M X$ is a k -space if and only if X is either discrete or locally compact and separable.

2. Proof of Theorem 1

First we shall describe a copy of the sequential hedgehog IS_ω in a topological group G admitting a functorial embedding of $[0, 1]$. Here the crucial role belongs to special trees consisting of closed intervals and ordered by the inverse inclusion relation, which will be called *usual Cantor schemas* throughout the paper.

We shall use the following notations: $\{0, 1\}^{<n} = \bigcup_{k < n} \{0, 1\}^k$, $\{0, 1\}^{\leq n} = \{0, 1\}^{<n+1}$, $\{0, 1\}^{<\omega} = \bigcup_{n \in \omega} \{0, 1\}^n$. In what follows we identify the unique element of $\{0, 1\}^0$ with the empty sequence \emptyset . For finite sequences $s = (s_0, \dots, s_n)$ and $t = (t_0, \dots, t_m)$ we denote by $s \hat{\ } t$ the *concatenation* of s and t , i.e., the finite sequence $(s_0, \dots, s_n, t_0, \dots, t_m)$. The sequence $(0, \dots, 0)$ of n zeros will be denoted by 0^n . In the same way we define the sequence 1^n . The length $l(s)$ of a sequence $s \in \{0, 1\}^{<\omega}$ is, by definition, the number $n \in \omega$ such that $s \in \{0, 1\}^n$.

Definition 4. *A family $\mathcal{J} = \{J_s : s \in \{0, 1\}^{<\omega}\}$ of subsets of \mathbb{R} is called a usual Cantor scheme, if it satisfies the following conditions:*

- (i) J_s is a closed subinterval of \mathbb{R} for all $s \in \{0, 1\}^{<\omega}$;
- (ii) $\min J_s = \min J_{s \hat{\ } 0} < \max J_{s \hat{\ } 0} < \min J_{s \hat{\ } 1} < \max J_{s \hat{\ } 1} = \max J_s$ for all $s \in \{0, 1\}^{<\omega}$; and
- (iii) the sequence $(\text{diam}(J_{(s_0, \dots, s_n)}))_{n \in \omega}$ converges to 0 for any $(s_i)_{i \in \omega} \in \{0, 1\}^\omega$.

For a usual Cantor scheme $\mathcal{J} = \{J_s : s \in \{0, 1\}^{<\omega}\}$ we make the following notations: $\mathcal{J}_n = \{J_s : s \in \{0, 1\}^n\}$, $J_s^{md} = [\max J_{s \hat{\ } 0}, \min J_{s \hat{\ } 1}]$ (here *md* comes from the word “middle”). A usual Cantor scheme \mathcal{J} is called *symmetric*, if the middle points of J_s and J_s^{md} coincide for all $s \in \{0, 1\}^{<\omega}$. In what follows all usual Cantor schemas are assumed to be symmetric.

For example, there is a canonical usual Cantor scheme $\mathcal{I} = \{I_s : s \in \{0, 1\}^{<\omega}\}$ appearing in the process of construction of the standard Cantor set $K \subset [0, 1]$ consisting of those numbers that have only 0's and 2's in their ternary expansion. Recall, that in order to obtain K we exclude from $[0, 1]$ open intervals step by step: $(\frac{1}{3}, \frac{2}{3})$ at the first step, $(\frac{1}{9}, \frac{2}{9})$ and $(\frac{7}{9}, \frac{8}{9})$ at the second step, and so on. Thus $I_{(0)} = [0, \frac{1}{3}]$, $I_{(1)} = [\frac{2}{3}, 1]$, $I_{(0,0)} = [0, \frac{1}{9}]$, \dots

Let $I \subset G$ be an embedding. For every $X = \{x_1, \dots, x_n\} \subset I$, $x_1 < \dots < x_n$, we set $\pi_-(X) = x_1^{-1} x_2 \cdots x_n^{(-1)^n}$, $\pi_+(X) = x_1 x_2^{-1} \cdots x_n^{(-1)^{n+1}}$, and $\pi_-(\emptyset) = \pi_+(\emptyset) = e$, where e is the neutral element of G . For a family \mathcal{A} of intervals we shall denote by $\partial \mathcal{A}$ the set $\bigcup_{I \in \mathcal{A}} \partial I$, where ∂I is the set of end-points of I . We shall also write $\text{diam}([a, b])$ for $|a - b|$.

¹In the rest of the paper the neutral element of a topological group G will be denoted by e_G or simply by e when G is clear from the context.

Given a usual Cantor scheme \mathcal{J} , we define for every $s \in \{0, 1\}^{<\omega}$ two maps $left_{s,\mathcal{J}}, right_{s,\mathcal{J}} : [0, 1] \rightarrow J_s$ such that $left_{s,\mathcal{J}}(0) = \min J_s$, $right_{s,\mathcal{J}}(0) = \max J_s$,

$$left_{s,\mathcal{J}}(1/3^n) = \max J_{s \frown 0^{n+1}} \text{ and } right_{s,\mathcal{J}}(1/3^n) = \min J_{s \frown 1^{n+1}}$$

for all $n \in \omega$, and $left_{s,\mathcal{J}}, right_{s,\mathcal{J}}$ are linear on every interval $[1/3^{n+1}, 1/3^n]$. It is clear that $left_{s,\mathcal{J}}, right_{s,\mathcal{J}}$ are continuous maps with the property

$$\lim_{\xi \rightarrow 0} left_{s,\mathcal{J}}(\xi) = \min J_s \text{ and } \lim_{\xi \rightarrow 0} right_{s,\mathcal{J}}(\xi) = \max J_s. \quad (1)$$

Whenever \mathcal{J} is clear from the context, we shall simply write $left_s$ and $right_s$ in place of $left_{s,\mathcal{J}}$ and $right_{s,\mathcal{J}}$. For every $n \in \omega$ and $\xi \in (0, 1]$ define the family $\mathcal{J}_{n,\xi}$ consisting of 2^{n+1} closed intervals as follows:

$$\mathcal{J}_{n,\xi} = \{left_s([0, \xi]), right_s([0, \xi]) : s \in \{0, 1\}^n\}$$

and define the map $z_{n,\mathcal{J}} : [0, 1] \rightarrow G$ letting $z_{n,\mathcal{J}}(0) = e$ and $z_{n,\mathcal{J}}(\xi) = \pi_-(\partial\mathcal{J}_{n,\xi})$ for all $\xi \in (0, 1]$. The continuity of maps $left_s, right_s$ implies the continuity of $z_{n,\mathcal{J}} : [0, 1] \rightarrow G$ for every $n \in \omega$. Moreover, it easily follows from equation (1) that $\lim_{\xi \rightarrow 0} z_{n,\mathcal{J}}(\xi) = e$, where e is the neutral element of G . Again, we shall write $z_n(\xi)$ instead of $z_{n,\mathcal{J}}(\xi)$ in case when \mathcal{J} is clear from the context.

Theorem 1 is a direct consequence of the following technical statement, which gives a description of a copy of the sequential hedgehog in a topological group admitting a functorial embedding of $[0, 1]$.

Proposition 5. *If \mathcal{J} is a usual Cantor scheme and $J_0 \subset G$ is a functorial embedding of the closed interval J_0 into a topological group G , then for every $n \in \omega$ the map $z_n : [0, 1] \rightarrow G$ is an embedding, and there exists a sequence $(d_n)_{n \in \omega} \in (0, 1]^\omega$ such that $\bigcup_{n \in \omega} z_n([0, d_n]) \subset G$ is a topological copy of the sequential hedgehog IS_ω .*

The proof of Proposition 5 will be split into a sequence of lemmas. The following is the most technically difficult among them.

Lemma 6. *If \mathcal{J} is a usual Cantor scheme and $J_0 \subset G$ is a functorial embedding, then $e \notin \bigcup_{n \in \omega} z_{n,\mathcal{J}}([d_n, 1])$ for every sequence $(d_n)_{n \in \omega} \in (0, 1]^\omega$.*

The proof of Lemma 6 will be also split into a sequence of simpler lemmas. The first of them is straightforward, and its proof is left to the reader. (We recall that $\mathcal{I} = \{I_s : s \in \{0, 1\}^{<\omega}\}$ is the canonical Cantor scheme.)

Lemma 7. *Let \mathcal{J} be a usual Cantor scheme. Then there exists an increasing homeomorphism $h : J_0 \rightarrow [0, 1]$ such that $h(J_s) = I_s$ for all $s \in \{0, 1\}^{<\omega}$.*

The following lemma easily follows from the compactness of $[0, 1]$.

Lemma 8. *Let $[0, 1] \subset G$ be an embedding and U be an open neighborhood of the neutral element $e \in G$. Then there exists $\varepsilon(U) > 0$ and an open neighborhood V of e such that for every $x_1, x_2 \in [0, 1]$ with $|x_1 - x_2| < \varepsilon(U)$ the following inclusion holds:*

$$\{x_1^{-1}Vx_2, x_1Vx_2^{-1}, x_2^{-1}Vx_1, x_2Vx_1^{-1}\} \subset U$$

For every $n > 0$ and $f : X \rightarrow X$ we shall denote by f^n the composition $\underbrace{f \circ \cdots \circ f}_n$. It will

be also convenient for us to denote by f^0 the identity map on X . Given any strictly increasing function $\varphi \in \omega^\omega$ such that $\varphi(0) > 0$ and a usual Cantor scheme \mathcal{J} , define a usual Cantor scheme $\tilde{\mathcal{J}} = \{\tilde{J}_s : s \in \{0, 1\}^{<\omega}\}$ letting $\tilde{J}_\emptyset = J_\emptyset$ and

$$\tilde{J}_{(s_0, \dots, s_n)} = J_{(s_0^{\varphi^2(0)} \wedge s_1^{\varphi^4(0) - \varphi^2(0)} \wedge \dots \wedge s_n^{\varphi^{2n+2}(0) - \varphi^{2n}(0)}}.$$

Thus $\tilde{\mathcal{J}}_m \subset \mathcal{J}_{\varphi^{2m}(0)}$ for all $m \in \omega$.

Claim 9. *Let $p \in \omega$ and $\varphi^{2p}(0) \leq n < \varphi^{2p+1}(0)$. Then $\partial \mathcal{J}_{n, \xi} \subset \{0, 1\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} \tilde{J}_s^{md}$ for all $\xi > 3^{n-\varphi(n)}$.*

Proof. Let us fix arbitrary $a \in \partial \mathcal{J}_{n, \xi}$ and $\xi > 3^{n-\varphi(n)}$. Assume, contrary to our claim, that a lies in the interior of \tilde{J}_s for some $s \in \{0, 1\}^{p+1}$. Concerning a , two cases are possible:

Case 1. $a \in \{left_t(0), right_t(0)\}$ for some $t \in \{0, 1\}^n$. Then $a \in \partial J_t$. Since $n < \varphi^{2p+2}(0)$ and $\tilde{J}_s \in \mathcal{J}_{\varphi^{2p+2}(0)}$, the inclusion $a \in \text{Int}(\tilde{J}_s)$ is impossible.

Case 2. There exists $t \in \{0, 1\}^n$ such that $a \in \{left_t(\xi), right_t(\xi)\}$. Without loss of generality, $a = left_t(\xi)$. Set $b = \min J_t = left_t(0)$. As it was already proven in Case 1, $b \in \{0, 1\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} \tilde{J}_s^{md}$. Again, two subcases are possible:

(i) b is an end-point of $\tilde{J}_{s_0}^{md}$ for some $s_0 \in \{0, 1\}^{\leq p}$, or $b = 0$. Then there exist $s_1 \in \{0, 1\}^p$ and $i \in \{0, 1\}$ such that $b \in \partial \tilde{J}_{s_1 \wedge i}$. Since $b = \min J_t$, we conclude that $i = 0$ and $b = \min \tilde{J}_{s_1 \wedge 0}$. Let $r \in \{0, 1\}^{\varphi^{2p}(0)}$ be such that $\tilde{J}_{s_1} = J_r$. The inequality $n \geq \varphi^{2p}(0)$ combined with $J_t \in \mathcal{J}_n$, $J_r \in \mathcal{J}_{\varphi^{2p}(0)}$, and $\min J_r = \min J_t = b$, implies that

$$a = left_t(\xi) \leq left_r(\xi) \leq left_r(1) = \max J_{r \wedge 0} < \min J_{r \wedge 1} \leq \min \tilde{J}_{s_1 \wedge 1}.$$

In addition,

$$a = left_t(\xi) > left_t(3^{n-\varphi(n)}) = \max J_{t \wedge 0^{\varphi(n)-n+1}} \geq \max J_{t \wedge 0^{\varphi^{2p+2}(0)-n}} = \max \tilde{J}_{s_1 \wedge 0}.$$

(The last equality immediately follows from $b = \min J_{t \wedge 0^{\varphi^{2p+2}(0)-n}} = \min \tilde{J}_{s_1 \wedge 0}$ and $J_{t \wedge 0^{\varphi^{2p+2}(0)-n}}, \tilde{J}_{s_1 \wedge 0} \in \mathcal{J}_{\varphi^{2p+2}(0)}$.) Therefore $a \in (\max \tilde{J}_{s_1 \wedge 0}, \min \tilde{J}_{s_1 \wedge 1}) = \text{Int}(\tilde{J}_{s_1}^{md})$, a contradiction.

(ii) $b \in \text{Int} \tilde{J}_{s_2}^{md}$ for some $s_2 \in \{0, 1\}^{\leq p}$. Again, let $r \in \{0, 1\}^{\varphi^{2p}(0)}$ be such that $\min J_r = \min J_t = b$. Since $\min J_r \in \text{Int}(\tilde{J}_{s_2}^{md})$, we conclude that $J_r \subset \text{Int}(\tilde{J}_{s_2}^{md})$. The inequality $n \geq \varphi^{2p}(0)$ combined with $\min J_r = \min J_t = b$ implies that $J_t \subset J_r \subset \text{Int}(\tilde{J}_{s_2}^{md})$. Therefore $a \in \text{Int}(\tilde{J}_{s_2}^{md})$ is an element of J_t , which contradicts the equation $a = left_t(\xi)$. \square

In the sequel the notation $\text{Auth}_+([a, b])$ stands for the family of all increasing autohomeomorphisms of the interval $[a, b]$.

Lemma 10. *Let φ and $\tilde{\mathcal{J}}$ be as above. Then for every indexed family $\{U_s : s \in \{0, 1\}^{<\omega}\}$ of open neighborhoods of the identity e of G there exists a sequence $(h_n)_{n \in \omega} \subset \text{Auth}_+([0, 1])$ such that*

- (1) $h_n(\tilde{\mathcal{J}})$ is a symmetric usual Cantor scheme;
- (2) $h_{n+1}|_{\bigcup_{s \in \{0, 1\}^{\leq n}} \tilde{J}_s^{md}} = h_n|_{\bigcup_{s \in \{0, 1\}^{\leq n}} \tilde{J}_s^{md}}$ for all $n \in \omega$; and

- (3) $\{\pi_-[h_n(\partial\mathcal{J}_{m,\xi} \cap \tilde{J}_s^{md})], \pi_+[h_n(\partial\mathcal{J}_{m,\xi} \cap \tilde{J}_s^{md})]\} \subset U_s$ for all $s \in \{0, 1\}^{\leq n}$, $m \geq \varphi^{2n}(0)$, and $\xi > 0$.

Proof. The following claim is the main building block of our proof. In order to shorten its proof, some explanations similar to those made in the proof of Claim 9 are omitted.

Claim 11. *Let \mathcal{J} , G , e be as in Lemma 10. Let also $s \in \{0, 1\}^{n_0}$, $m > 0$, $c = \max J_{s \wedge 0^m}$, $d = \min J_{s \wedge 1^m}$, and let U be an open neighborhood of e in G . Then there exists $h \in \text{Auth}_+(J_s)$ such that*

- (i) $h|_{J_{s \wedge 0^m}}$ and $h|_{J_{s \wedge 1^m}}$ are linear;
- (ii) $\text{diam}(h(J_{s \wedge 0^m})) = \text{diam}(h(J_{s \wedge 1^m}))$; and
- (iii) $\{\pi_-(h(\partial\mathcal{J}_{n,\xi} \cap [c, d])), \pi_+(h(\partial\mathcal{J}_{n,\xi} \cap [c, d]))\} \subset U$ for all $n \geq n_0$ and $\xi \in (0, 1]$.

Proof. Let us find some $a, b \in J_s$ such that $a < b$, the middle points of $[a, b]$ and J_s coincide, and $|a - b| < \varepsilon(U)$. The latter means that there exists an open neighborhood V of e such that $u^{-1}Vv \cup uVv^{-1} \subset U$ for all $u, v \in [a, b]$. The continuity of the group operation on G gives us a sequence $(V_n)_{n \in \omega}$ of open neighborhoods of e such that $V_n^{2^n} \subset V$. Let $h \in \text{Auth}_+(J_s)$ be such that the following conditions are satisfied: $h|_{[\min J_s, c]}$ and $h|_{[d, \max J_s]}$ are linear bijections onto $[\min J_s, a]$ and $[b, \max J_s]$ respectively, and $\text{diam}(h(J_t)) < \varepsilon(V_{l(t)})$ provided $J_t \subset (c, d)$. The existence of h follows from Lemma 7. Given arbitrary $n \geq n_0$ and $\xi \in [0, 1]$, write the family $\{J \in \mathcal{J}_{n,\xi} : J \subset [c, d]\}$ in the form $\{J_1, \dots, J_q\}$ such that J_i is situated to the left of J_j provided $i < j$. Let us note that each $J \in \mathcal{J}_{n,\xi}$ is contained in some element of \mathcal{J}_{n+1} , and hence $q \leq 2^{n+1}$. It can be easily derived from the definitions of a usual Cantor scheme and maps $left_-, right_-$ that $\{c, d\} \subset \partial\mathcal{J}_{n,\xi}$ for all n, ξ such that $\xi = 3^{-(n_0+m-n-1)}$ or $n \geq n_0 + m$, and $\{c, d\} \cap \partial\mathcal{J}_{n,p} = \emptyset$ otherwise. In the first case we have

$$\pi_-[h(\partial\mathcal{J}_{n,\xi} \cap [c, d])] = a^{-1}\pi_+(h(\partial J_1))\pi_+(h(\partial J_2)) \cdots \pi_+(h(\partial J_q))b \subset a^{-1}V_{n+1}^q b \subset a^{-1}Vb \subset U.$$

In the second case $n < n_0 + m$ and $\xi \neq 3^{-(n_0+m-n-1)}$. If $\xi < 3^{-(n_0+m-n-1)}$, then (c, d) contains all elements of $\mathcal{J}_{n,\xi}$ whose intersection with $[c, d]$ is nonempty, and hence

$$\pi_-[h(\partial\mathcal{J}_{n,\xi} \cap [c, d])] = \pi_-(h(\partial J_1))\pi_-(h(\partial J_2)) \cdots \pi_-(h(\partial J_q)) \subset V_{n+1}^q \subset U.$$

It suffices to consider the case $\xi > 3^{-(n_0+m-n-1)}$. Let $u = left_s(\xi)$ and $v = right_s(\xi)$. Then $c < u < v < d$ and $J_i \subset (u, v)$ for all $i \leq q$. Therefore

$$\pi_-[h(\partial\mathcal{J}_{n,\xi} \cap [c, d])] = u^{-1}\pi_+(h(\partial J_1))\pi_+(h(\partial J_2)) \cdots \pi_+(h(\partial J_q))v \subset u^{-1}V_{n+1}^q v \subset u^{-1}Vv \subset U.$$

Verification that $\pi_+(h(\partial\mathcal{J}_{n,\xi} \cap [c, d])) \subset U$ is similar, and thus condition (iii) is satisfied. \square

Applying Claim 11 for the usual Cantor scheme \mathcal{J} , $s = \emptyset$, $m = \varphi^2(0)$, and $U = U_\emptyset$, we get $h_0 \in \text{Auth}_+(\tilde{J}_\emptyset)$ satisfying the conditions (i) – (iii) of Claim 11. Conditions (i) and (ii) obviously imply (1), condition (iii) implies (3), while (2) is trivial.

Assuming that we have already constructed h_k satisfying (1) – (3) for all $k < n$, set $h_n|_{\bigcup_{s \in \{0,1\}^{\leq n-1}} \tilde{J}_s^{md}} = h_{n-1}|_{\bigcup_{s \in \{0,1\}^{\leq n-1}} \tilde{J}_s^{md}}$. Thus condition (2) is satisfied. In addition, (3) holds for all $s \in \{0, 1\}^{\leq n-1}$. It suffices to define h_n on $[0, 1] \setminus \bigcup_{s \in \{0,1\}^{\leq n-1}} \tilde{J}_s^{md} = \bigcup_{s \in \{0,1\}^n} (\tilde{J}_s \setminus \partial\tilde{J}_s)$. Let us note, that for every particular $s \in \{0, 1\}^n$ the construction of $h_n|_{\tilde{J}_s}$ is similar to that of h_0 . Given any $s = (s_0, \dots, s_{n-1}) \in \{0, 1\}^n$, set $t = s_0^{\varphi^2(0)} \wedge s_1^{\varphi^4(0) - \varphi^2(0)} \wedge \dots \wedge s_{n-1}^{\varphi^{2n}(0) - \varphi^{2n-2}(0)}$

and $m = \varphi^{2n+2}(0) - \varphi^{2n}(0)$. Thus $J_t = \tilde{J}_s$. Applying Claim 11 for \mathcal{J} , $t \in \{0, 1\}^{\varphi^{2n}(0)}$, m , and U_s , we get $h_s \in \text{Auth}_+(J_t)$ satisfying conditions (i) – (iii) of Claim 11. Set $h_n|_{\tilde{J}_s} = h_s \circ h_{n-1}|_{\tilde{J}_s}$. Again, (i) and (ii) imply (1), and (iii) implies (3). \square

The following simple statement is due to M. Tkachenko (see [17, Lemma 1.3]).

Lemma 12. *Let G be a topological group and \mathcal{U} be the family of all open neighborhood of the neutral element of G . Then for every $U \in \mathcal{U}$ there exists a decreasing sequence $(U_n)_{n \in \omega} \subset \mathcal{U}$ such that $U_{\sigma(0)}U_{\sigma(1)} \cdots U_{\sigma(n)} \subset U$ for every bijection $\sigma : \{0, \dots, n\} \rightarrow \{0, \dots, n\}$.*

The proof of the following simple technical lemma is left to the reader.

Lemma 13. *For every function $\varphi : \omega \rightarrow \omega \setminus \{0\}$ there exist strictly increasing functions $\psi, \theta : \omega \rightarrow \omega$ such that*

(i) $\varphi(k) < \psi(k) < \theta(k)$ for all $k \in \omega$; and

(ii) $\psi^{n+1}(0) = \theta^n(0)$ for all $n \in \omega$.

Proof of Lemma 6. Throughout the proof we denote by $\hat{h} : G \rightarrow G$ some continuous homomorphism extending $h \in \text{Auth}_+(J_\emptyset)$. Let \mathcal{U} be the family of all open neighborhoods of the neutral element e of G . The continuity of the group operation of G yields the existence of an element $U \in \mathcal{U}$ such that $U \cap 0^{-1}U1 = \emptyset$. By Lemma 12 there exists a sequence $(U_n)_{n \in \omega} \subset \mathcal{U}$ such that $U_{i_1}U_{i_2} \cdots U_{i_{2^{n+1}-1}} \subset U$ for any $n \in \omega$ and $(i_1, \dots, i_{2^{n+1}-1})$ such that $|\{j < 2^{n+1} : i_j = k\}| = 2^k$ for all $k \in \{0, \dots, n\}$.

Let $\varphi : \omega \rightarrow \omega$ be a strictly increasing map with the property $3^{n-\varphi(n)} < d_n$ for every $n \in \omega$. Let us fix a sequence $(h_n)_{n \in \omega} \subset \text{Auth}_+([0, 1])$ satisfying conditions (1) – (3) of Lemma 10 with U_s equal to $U_{l(s)}$ defined above, where $s \in \{0, 1\}^{<\omega}$. Conditions (1), (2) imply that $|h_n(t) - h_{n+1}(t)| \leq 2^{-n-1}$, and hence the sequence $(h_n)_{n \in \omega}$ is uniformly convergent to a monotone continuous surjective function $h : [0, 1] \rightarrow [0, 1]$. For every $s \in \{0, 1\}^{\leq n}$ we have $h|_{\tilde{J}_s^{md}} = h_n|_{\tilde{J}_s^{md}}$ by (2), consequently $h|_{\tilde{J}_s^{md}}$ is not constant for every $s \in \{0, 1\}^{<\omega}$. Since $\bigcup_{s \in \{0, 1\}^{<\omega}} \tilde{J}_s^{md}$ is dense in $[0, 1]$, we conclude that $h \in \text{Auth}_+([0, 1])$ is a monotone continuous surjection which fails to be constant on arbitrary open subset of $[0, 1]$. We claim that

$$\hat{h}\left(\bigcup_{p \in \omega} \bigcup_{\varphi^{2p}(0) \leq n < \varphi^{2p+1}(0)} \{z_{n, \mathcal{J}}(\xi) : \xi > 3^{n-\varphi(n)}\}\right) \subset 0^{-1}U1.$$

Indeed, let us fix arbitrary $p \in \omega$, $\varphi^{2p}(0) \leq n < \varphi^{2p+1}(0)$, and $\xi \in (3^{n-\varphi(n)}, 1]$. Then by Claim 9 we have $\partial \mathcal{J}_{n, \xi} \subset \{0, 1\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} \tilde{J}_s^{md}$. Therefore

$$z_{n, \mathcal{J}}(\xi) = \pi_-(\partial \mathcal{J}_{n, \xi}) = \pi_-(\{0\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} (\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_s^{md}) \cup \{1\}).$$

Combining the equation above with (2) and (3) of Lemma 10, we conclude that

$$\begin{aligned} \hat{h}(z_{n, \mathcal{J}}(\xi)) &= \hat{h}[\pi_-(\{0\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} (\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_s^{md}) \cup \{1\})] = \\ &= \pi_-[h(\{0\} \cup \bigcup_{s \in \{0, 1\}^{\leq p}} (\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_s^{md}) \cup \{1\})] = \\ &= \pi_-(\{0\} \cup h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_1}^{md}) \cup h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_2}^{md}) \cup \cdots \cup h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_{2^{p+1}-1}}^{md}) \cup \{1\}) = \\ &= 0^{-1} \pi_{\delta_1}(h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_1}^{md})) \pi_{\delta_2}(h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_2}^{md})) \cdots \pi_{\delta_{2^{p+1}-1}}(h_p(\partial \mathcal{J}_{n, \xi} \cap \tilde{J}_{s_{2^{p+1}-1}}^{md})) 1 \subset \\ &\subset 0^{-1} U_{l(s_1)} U_{l(s_2)} \cdots U_{l(s_{2^{p+1}-1})} 1 \subset 0^{-1} U 1, \end{aligned}$$

where $\{s_1, \dots, s_{2^{p+1}-1}\}$ is the enumeration of $\{0, 1\}^{\leq p}$ such that $\tilde{I}_{s_i}^{md}$ is situated to the left of $\tilde{I}_{s_j}^{md}$ provided $1 \leq i < j < 2^{p+1}$, and $\delta_i \in \{+, -\}$. Then $\bigcup_{p \in \omega} \bigcup_{\varphi^{2p}(0) \leq n < \varphi^{2p+1}(0)} \{z_{n, \mathcal{J}}(\xi) : 1 \geq \xi > 3^{n-\varphi(n)}\} \cap \hat{h}^{-1}(U) = \emptyset$ by our choice of U .

Let $\psi, \theta : \omega \rightarrow \omega$ be increasing number sequences such as in Lemma 13, i.e. $\varphi(n) \leq \psi(n) \leq \theta(n)$ for all $n \in \omega$ and $\theta^n(0) = \psi^{n+1}(0)$ for all $n \geq 1$. It follows from the above that there are continuous homomorphisms $h_\psi, h_\theta : G \rightarrow G$ such that

$$\begin{aligned} \bigcup_{p \in \omega} \bigcup_{\psi^{2p}(0) \leq n < \psi^{2p+1}(0)} \{z_{n, \mathcal{J}}(\xi) : 1 \geq \xi > 3^{n-\psi(n)}\} \cap h_\psi^{-1}(U) &= \emptyset \text{ and} \\ \bigcup_{p \in \omega} \bigcup_{\theta^{2p}(0) \leq n < \theta^{2p+1}(0)} \{z_{n, \mathcal{J}}(\xi) : 1 \geq \xi > 3^{n-\theta(n)}\} \cap h_\theta^{-1}(U) &= \emptyset, \end{aligned}$$

which implies $\bigcup_{n \in \omega} \{z_{n, \mathcal{J}}(\xi) : 1 \geq \xi > 3^{n-\varphi(n)}\} \cap (h_\psi^{-1}(U) \cap h_\theta^{-1}(U)) = \emptyset$, and hence the fact that $e \notin \overline{\bigcup_{n \in \omega} \{z_{n, \mathcal{J}}(\xi) : 1 \geq \xi > d_n\}}$ is proven. \square

For a subset X of a topological group G we shall denote by $\langle X \rangle$ the smallest subgroup of G containing X .

Lemma 14. *Let $[0, 1] \subset G$ be a functorial embedding, $\{x_i : 0 \leq i \leq n\} \cup \{y_j : 0 \leq j \leq m\} \subset [0, 1]$, and $y_{j_0} \notin \{y_j : j \neq j_0\} \cup \{x_i : 0 \leq i \leq n\}$ for some j_0 . Then*

$$x_0^{k_0} \cdot x_1^{k_1} \cdot \dots \cdot x_n^{k_n} \neq y_0^{l_0} \cdot y_1^{l_1} \cdot \dots \cdot y_m^{l_m}$$

for arbitrary integers k_i, l_i such that $l_{j_0} \in \{-1, 1\}$.

In particular, for every $n \in \omega$ and usual Cantor scheme \mathcal{J} the map $z_{n, \mathcal{J}} : [0, 1] \rightarrow G$ is an embedding, and $z_{n, \mathcal{J}}((0, 1]) \cap z_{m, \mathcal{J}}((0, 1]) = \emptyset$ for all $m \neq n$.

Proof. The second statement easily follows from the first one. To prove the first statement, set $x = x_0^{k_0} \cdot x_1^{k_1} \cdot \dots \cdot x_n^{k_n}$, $y = y_0^{l_0} \cdot y_1^{l_1} \cdot \dots \cdot y_m^{l_m}$, and assume to the contrary that $x = y$. Let $h \in \text{Auth}_+([0, 1])$ be such that $h(u) = u$ for all $u \in \{y_j : j \neq j_0\} \cup \{x_i : 0 \leq i \leq n\}$ and $h(y_{j_0}) \neq y_{j_0}$. Since the embedding $[0, 1] \subset G$ is functorial, there exists a continuous homomorphism $\hat{h} : G \rightarrow G$ extending h . It follows from the above that $\hat{h}(x) = x = y$ and $\hat{h}(y) \neq y$, a contradiction. \square

Proof of Proposition 5. In light of Lemmas 6 and 14 we are left with the task of constructing a sequence $(d_n)_{n \in \omega}$ of positive reals with the property $z_n((0, d_n]) \cap \overline{\bigcup_{k > n} z_k([0, d_k])} = \emptyset$ for all $n \in \omega$. Let $A = \{\xi \in (0, 1] : z_0(\xi) \in \overline{\bigcup_{k > 0} z_k([0, 1])}\}$. Since $\lim_{\xi \rightarrow 0} z_k(\xi) = e$ for all $k \in \omega$, we conclude that A consists of $\xi \in [0, 1]$ such that $\xi \in \overline{\bigcup_{k > 0} z_k([c_k, 0])}$ for some sequence $(c_k)_{k > 0}$ of positive reals.

We claim that $0 \notin \overline{A}$. Indeed, assuming the converse we could find a sequence $(\xi_n)_{n \in \omega}$ of elements of A converging to 0. It follows from the above that for every $n \in \omega$ there exists a sequence $(c_{n,k})_{k > 0}$ of positive reals such that $z_0(\xi_n) \in \overline{\bigcup_{k > 0} z_k([c_{n,k}, 1])}$. Set $c_k = \min\{c_{n,k} : n \leq k\}$. Then $e \in \overline{\bigcup_{k > 0} z_k([c_k, 1])}$, which contradicts Lemma 6.

It follows from the above that $A \subset (d_0, 1]$ for some $d_0 > 0$, and consequently $z_0((0, d_0]) \cap \overline{\bigcup_{k > 0} z_k([0, 1])} = \emptyset$. In the same way for every $n > 0$ we can find d_n such that $z_n((0, d_n]) \cap \overline{\bigcup_{k > n} z_k([0, 1])} = \emptyset$, which completes our proof. \square

Remark. Let \mathcal{J} be a usual Cantor scheme such that $J_\emptyset \subset G$ is a functorial embedding. Let also $C_n = z_{n,\mathcal{J}}([0, d_n])$ for a sequence $(d_n)_{n \in \omega}$ fulfilling the requirements of Proposition 5. Then it can be easily derived from Lemma 14 that the map

$$\prod_{i \leq n} [0, d_n] \ni (\xi_0, \dots, \xi_n) \mapsto \prod_{i \leq n} z_{n,\mathcal{J}}(\xi_i)$$

is a homeomorphism, and we denote its image by D_n . Thus we have an increasing sequence

$$D_0 \subset D_1 \subset \dots \subset D_n \subset \dots,$$

where each D_n is a homeomorphic copy of the $(n+1)$ -dimensional cube². Let ν be the topology of G and let τ be the strongest topology on $D = \bigcup_{n \in \omega} D_n$ such that $\tau|_{D_n} = \nu|_{D_n}$ for all $n \in \omega$. It is easy to see that (D, τ) contains a homeomorphic copy of $\mathbb{R}^\infty = \lim_{\leftarrow} \mathbb{R}^n$, which is homeomorphic to the Markov free topological group over $[0, 1]$ (see [19]). But we do not know whether $\nu|_D = \tau|_D$. This leads us to the following question.

Question 15. *Let $[0, 1] \subset G$ be a functorial embedding. Does G contain a topological copy of the Markov free topological group over $[0, 1]$? More precisely, does the construction described above yield a copy of \mathbb{R}^∞ in G ?*

Similarly to the proof of Proposition 16 below, the positive solution of the above question would imply that for every functor F of a free topological group and every Tychonov space X containing a topological copy of $[0, 1]$, the group FX contains a topological copy of the Markov free topological group over $[0, 1]$. \square

3. Proof of Theorem 2

We shall derive Theorem 2 from the following slightly more general statement, where we specify the properties of the category \mathcal{T} used in the proof of Theorem 2.

Proposition 16. *Let (F, i) be a functor of a free topological group on a category \mathcal{T} of topological spaces such that $I \in \text{Ob}(\mathcal{T})$ and $\text{Mor}(I, I) \supset \text{Auth}_+([0, 1])$. The group FX over an object X of \mathcal{T} contains:*

- (1) *a copy of the Fréchet-Urysohn fan S_ω provided there is a morphism $f : X \rightarrow I$ in \mathcal{T} whose restriction $f|_Q$ onto some subspace $Q \subset X$ without isolated points is an embedding of Q into I , and $\text{Mor}(I, I)$ contains all continuous maps $\phi : I \rightarrow I$; and*
- (2) *a copy of the sequential hedgehog IS_ω provided there is a copy $Y \subset X$ of I , a surjective map $f \in \text{Mor}(X, Y)$, and a homeomorphism $h \in \text{Mor}(Y, I)$.*

Proposition 16 is a consequence of Theorem 1 and the following

Lemma 17. *Let X be a Tychonov space containing a topological copy Q of the space of rational numbers and \mathcal{J} be a usual Cantor scheme with $J_\emptyset = [0, 1]$. Then there exists a homeomorphic copy $Q' \subset Q$ of Q , and a continuous map $\psi : X \rightarrow J_\emptyset$ such that $\psi|_{Q'}$ is a homeomorphism between the spaces Q' and $\partial\mathcal{J} = \bigcup_{s \in \{0,1\}^{<\omega}} \partial J_s$.*

²This gives an alternative (but much longer) proof of the fact [4] that each topological group is infinite-dimensional provided it admits a functorial embedding of the closed interval.

Proof. Let $I = [0, 1]$. It is well-known that the diagonal map $\delta : X \rightarrow I^{\text{Mor}(X, I)}$ is an embedding, where $\text{Mor}(X, I)$ stands for the set of all continuous maps from X to I . Assume that X contains a subset $Q \subset X$ homeomorphic to the space of rational numbers. Using the fact that $\delta(Q) \subset I^{\text{Mor}(X, I)}$ has a countable base, one can construct a countable subset $C \subset \text{Mor}(X, I)$ such that the restriction $\text{pr}|\delta(Q)$ of the projection $\text{pr} : I^{\text{Mor}(X, I)} \rightarrow I^C$ is a homeomorphic embedding. By the standard argument (see [10, Theorem 21.18]), we can find a topological copy $Q_1 \subset \text{pr} \circ \delta(Q)$ of \mathbb{Q} whose closure $\overline{Q_1}$ in the (metrizable) cube I^C is zero-dimensional, and hence is homeomorphic to the Cantor space $\{0, 1\}^\omega$. Let

$$e : \overline{Q_1} \rightarrow \bigcup_{(s_n)_{n \in \omega} \in \{0, 1\}^\omega} \bigcap_{n \in \omega} J_{(s_0, \dots, s_n)}$$

be the homeomorphism with the property $e(Q_1) = \partial\mathcal{J}$ (its existence follows from [5, Part 4, Th. 1] or the main result of [9]), and $\bar{e} : I^C \rightarrow I$ be an extension of e to a continuous map. Then the set $Q' = Q \cap (\text{pr} \circ \delta)^{-1}(Q_1)$ is a topological copy of \mathbb{Q} and the map $f = \bar{e} \circ \text{pr} \circ \delta : X \rightarrow I$ restricted to Q' is a homeomorphism between Q' and $\partial\mathcal{J}$. \square

Observe that for an arbitrary family $\{(x_k^n)_{k \in \omega} : n \in \omega\}$ of sequences of elements of $(0, 1]$, the subspace $\{x_k^n \times \{n\} : n, k \in \omega\} \cup \{\{0\} \times \omega\}$ of IS_ω is homeomorphic to S_ω provided $\lim_{k \rightarrow \infty} x_k^n = 0$ for all $n \in \omega$.

Proof of Proposition 16. Throughout the proof we shall identify X with $i_X(X)$. We present here only the proof of the first part. The proof of the second one is analogous.

Let $f \in \text{Mor}(X, I)$ be such that $f|Q$ is an embedding for some homeomorphic copy $Q \subset X$ of \mathbb{Q} . Lemma 17 implies that there exists a usual Cantor scheme \mathcal{J} , a continuous map $g : I \rightarrow I$, and a copy $Q_1 \subset f(Q)$ of \mathbb{Q} such that $g(Q_1) = \partial\mathcal{J}$ and $g|Q_1$ is an embedding. Then $h = g \circ f$ belongs to $\text{Mor}(X, I)$ and $h(f^{-1}(Q_1)) = \partial\mathcal{J}$. Set $r_{n,k} = z_{n,\mathcal{J}}(3^{-k})$. Applying Proposition 5, we can find a sequence $(k_n)_{n \in \omega}$ of natural numbers such that $R = \{e\} \cup \{r_{n,k} : n \in \omega, k \geq k_n\}$ is a homeomorphic copy of S_ω . By our construction of the maps $z_{n,\mathcal{J}}$, for every $n, k \in \omega$ we can find elements

$$0 = u_{n,k,0} < u_{n,k,1} < \dots < u_{n,k,2^{n+2}-1} = 1$$

of $\partial\mathcal{J}$ such that $r_{n,k} = u_{n,k,0}^{-1} u_{n,k,1} u_{n,k,2}^{-1} \dots u_{n,k,2^{n+2}-1}$. In addition, $u_{n,k,p}$ does not depend on k provided 4 divides p or $p+1$, and

$$\lim_{k \rightarrow \infty} u_{n,k,4q+1} = u_{n,k,4q}, \quad \lim_{k \rightarrow \infty} u_{n,k,4q+2} = u_{n,k,4q+3} \quad (2)$$

for all $q < 2^n - 1$. Set $C = (f|Q)^{-1}(Q_1)$, $v_{n,k,p} = (h|C)^{-1}(u_{n,k,p})$,

$$y_{n,k} = v_{n,k,0}^{-1} v_{n,k,1} v_{n,k,2}^{-1} \dots v_{n,k,2^{n+2}-1},$$

and $Y = \{e_{FX}\} \cup \{y_{n,k} : n \in \omega, k \geq k_n\}$. Since $h|C : C \rightarrow \partial\mathcal{J}$ is a homeomorphism, the sequence $(y_{n,k})_{k \in \omega}$ converges to e_{FX} for all $n \in \omega$ by (2). In addition, the continuous homomorphism $Fh : FX \rightarrow FI$ maps $y_{n,k}$ to $r_{n,k}$ by our choice of $v_{n,k,p}$, and hence $Fh(Y) = R$. By the definition, S_ω is the union of a countable family of disjoint convergent sequences whose limit points coincide endowed with the strongest topology in which these sequences are still convergent. Thus the continuity of $Fh|Y$ implies that Y is homeomorphic to R . \square

Proof of Theorem 2. Follows immediately from Proposition 16, Lemma 17, and the fact that $[0, 1]$ is an absolute retract in the category of Tychonov spaces. \square

Lemma 18. *Under the assumptions of Corollary 1 the image $i_X(X)$ is closed in FX .*

Proof. Throughout the proof we shall identify $Z \in \text{Ob}(\mathcal{T})$ with $i_Z(Z)$. Let $j : X \rightarrow \bar{X}$ be the inclusion. It suffices to show that $X = (Fj)^{-1}(\bar{X})$. Assuming the converse, by the minimality of F we could find a finite subset $\{x_i : i \leq n\} \subset X$ and integers $m_i, i \leq n$, such that $x^* := (Fj)(x_0^{m_0} \cdots x_n^{m_n}) \in \bar{X} \setminus X$. Therefore the elements x^* and $x_0^{m_0} \cdots x_n^{m_n}$ of $F\bar{X}$ coincide. Let $f : \bar{X} \rightarrow [0, 1]$ be a continuous map such that $f(x^*) \neq f(x_i)$ for all $i \leq n$. From the above it follows that

$$f(x^*) = Ff(x^*) = Ff(x_0^{m_0} \cdots x_n^{m_n}) = f(x_0)^{m_0} \cdots f(x_n)^{m_n},$$

which contradicts Lemma 14. □

Proof of Corollary 1. Let us denote by \mathbb{L} the subspace $\{(0, 0)\} \cup \{(\frac{1}{n}, \frac{1}{mn}) : n, m > 0\}$ of \mathbb{R}^2 . According to [6, Lemma 8.3], a first countable space contains a closed topological copy of \mathbb{L} if and only if it fails to be locally compact.

Assume, contrary to our claim, that X is not scattered and fails to be locally compact. Then X contains a topological copy of the space \mathbb{Q} as well as a closed topological copy of \mathbb{L} . Since FX is generated by its second-countable subspace X , it has countable pseudocharacter. In addition, it is normal being Lindelöf and Tychonov. Applying Theorem 2, we conclude that FX contains a topological copy of S_ω . It is well-known [13] that a topological group G contains a (closed) topological copy of S_ω if and only if it contains a (closed) topological copy of the Arens' space S_2 , see [13] for its definition. It was shown by Lin [12, Corollary 2.6] that a regular space Z with countable pseudocharacter contains a topological copy of S_ω if and only if it contains a closed topological copy of S_ω . It follows from the above that FX contains a closed topological copy of S_ω . In addition, FX contains a closed copy of \mathbb{L} by Lemma 18, which contradicts [3, Theorem 1] asserting that a normal k -space containing closed topological copies of \mathbb{L} and S_ω is not homeomorphic to any topological group. □

References

- [1] A. V. Arkhangel'skiĭ, *Frequency spectrum of a topological space and classification of spaces*, Dokl. Akad. Nauk SSSR 206 (1972), 265–268. (in Russian)
- [2] A. V. Arkhangel'skiĭ, O. G. Okunev, V. G. Pestov, *Free topological groups over metrizable spaces*, Topology Appl. **33** (1989), 63–76.
- [3] T. O. Banach, *On topological groups containing a Fréchet-Urysohn fan*, Matematychni Studii **9** (1998), 145–154.
- [4] T. O. Banach, M. M. Zarichnyĭ, *The interval $[0, 1]$ admits no functorial embedding into a finite-dimensional or metrizable topological group*, Serdica Math. J. **26** (2000), 1–4.
- [5] C. Bessaga, A. Pelczyński, *Selected Topics in Infinite-Dimensional Topology*, PWN, Warszawa, 1975.
- [6] E. K. van Douwen, *The integers and topology*, Handbook of Set-Theoretic Topology, K. Kunen and J. E. Vaughan, Eds., North-Holland, Amsterdam, 1984, pp. 111–167.

- [7] K. H. Hofmann, S. A. Morris, *Free compact groups I: free compact abelian groups*, Topology Appl. **23** (1986), 41–64.
- [8] K. H. Hofmann, S. A. Morris, *Free compact groups IV: Splitting the component and the structure of the commutator group*, J. Pure Appl. Algebra **70** (1991), 89–96.
- [9] M. Hrušak, B. Zamora Aviles, *Countable dense homogeneity of definable spaces*, Proc. Amer. Math. Soc. **133**:11 (2005), 3429–3435.
- [10] A. Kechris, *Classical Descriptive Set Theory*, Grad. Texts Math. **156**, Springer-Verlag, Berlin, 1995.
- [11] J. Kulesza, *Spaces which do not embed in topological groups of the same dimension*, Topology Appl. **50** (1993), 139–145.
- [12] S. Lin, *A note on Arens space and sequential fan*, Topology Appl. **81** (1997), 185–197.
- [13] J. Nagata. *Generalized Metric Spaces, I.*, Topics in General Topology, K.Morita, J.Nagata, Eds., North-Holland, Amsterdam, 1989), pp. 315–366.
- [14] P. Nickolas, M. Tkachenko, *The character of free topological groups, I*, Appl. Gen. Top. **6**:1 (2005), 15–41.
- [15] P. Nickolas, M. Tkachenko, *The character of free topological groups, II*, Appl. Gen. Top. **6**:1 (2005), 43–56.
- [16] D. B. Shakhmatov, *A categorial version of the Lefschetz–Nöbeling–Pontryagin theorem on embedding compacta in \mathbb{R}^n* , Topology Appl. **85** (1998), 345–349.
- [17] M. G. Tkačenko, *On topologies of free groups*, Czechoslovak Math. J. **34**:109 (1984), 541–551.
- [18] J. E. Vaughan, *Small uncountable cardinals and topology*, Open problems in topology, J. van Mill, G.M. Reed, Eds., North-Holland, Amsterdam, 1990, pp. 197–216.
- [19] M. M. Zarichnyi, *Free topological groups of absolute neighborhood retracts and infinite-dimensional manifolds*, Dokl. Akad. Nauk SSSR **266**:3 (1982), 541–544. (in Russian)

Taras Banakh, Instytut Matematyki, Akademia Świętokrzyska, Kielce, Poland and Department of Mechanics and Mathematics, Ivan Franko Lviv National University, Universytetska 1, Lviv, 79000, Ukraine.

E-mail address: tbanakh@franko.lviv.ua

Dušan Repovš, Institute of Mathematics, Physics and Mechanics, and Faculty of Education, University of Ljubljana, Jadranska 19, P.O.B. 2964, Ljubljana, Slovenija 1001.

E-mail address: dusan.repovs@guest.arnes.si

Lyubomyr Zdomsky, Department of Mechanics and Mathematics, Ivan Franko Lviv National University, Universytetska 1, Lviv, 79000, Ukraine.

E-mail address: lzdomsky@gmail.com